

## Fully reduced-HMGB1, LPS-free

**Product Number:** \*\*\*\*\*

**Expiration date:** \*\*\*\*\*

**Batch number:** \*\*\*\*\*

**Batch concentration:** \*\*\*\*\* mg/mL after addition of  
\*\*\*\*\*  $\mu$ L of distilled water.

### Product Description:

HMGB1 is a 25 kDa nuclear protein, present in almost all mammalian cells. The protein is virtually identical (213/215 aa) in human, mouse, rat. This product corresponds to the human sequence and is produced in *E.coli*. Fully reduced-HMGB1 (complete notation: HMGB1C23hC45hC106h - Antoine J. *et al* (2014).Mol Med) forms complex with CXCL12 and has chemoattractant activity. It DOES NOT induce cytokine/chemokine secretion when given to target cells. The protein is free from LPS (<0.1EU/mL).

The product contains <0.006% v/v of Triton X-114 due to LPS removal procedure.

### Reagent format:

**Fully reduced-HMGB1 we provide is the natural protein, with no tags or additional amino acids.**

The lyophilized protein once reconstituted will be dissolved in a solution containing 50 mM HEPES pH 7.9, 500 mM NaCl, DTT 0,5 mM.

**Storage:** 2-8°C. The protein once resuspended can be stored frozen (-20°C).

Oxidation of cysteine 106 makes the protein inactive (Kazama *et al*, Immunity 2008; 29, 21-32).

To avoid cysteine oxidation DTT 0.5 mM is added during protein purification.

### How to use product:

The product can be used in cell migration assays, both *in vitro* and *in vivo*; maximum activity is at 1 nM (Palumbo *et al*, 2004). Intraperitoneal injection in the mouse recruits neutrophils, monocytes and macrophages (Penzo *et al*, 2010).

### This product is for research use only

#### References:

- Agnieszka I. *et al* (2021) Antioxidants The Time-Course of Antioxidant Irisin Activity: Role of the Nrf2/HO-1/HMGB1 Axis 10(1):88
- Li R. *et al* (2021) Lipopolysaccharide-Activated Canine Platelets Upregulate High Mobility Group Box-1 via Toll-Like Receptor. *Front Vet Sci.* 8:674678
- Piao C. *et.al* (2020) An 8-Hydroxy-Quinoline Derivative Protects Against Lipopolysaccharide-Induced Lethality in Endotoxemia by Inhibiting HMGB1-Mediated Caspase-11 Signaling. *Frontiers in Pharmacology.* 12:673818
- Piao C. *et al* (2020) A RAGE-antagonist peptide potentiates polymeric micelle-mediated intracellular delivery of plasmid DNA for acute lung injury gene therapy. 12(25):13606-13617

```

MGKGDPPKPR  GKMSYAFFV  QTCREEHKKK
HPDASVNFSE  FSKKCSERWK  TMSAKEKGF
EDMAKADKAR  YEREMKTYIP  PKGETKKKFK
DPNAPKRPPS  AFFLFCSEYR  PKIKGEHPGL
SIGDVAKKLG  EMWNTAADD  KQPYEKKA
LKEKYEKIDIA  AYRAKGKPPA  AKKGVVKA
SKKKKEEEDD  EEEDEDEEEE  EEEDEDEEEE
DDDDE
    
```

Fig. 1. Fully reduced-HMGB1 sequence




Fig. 2. SDS-PAGE with Coomassie Blue staining




Fig. 3. Reconstructed molecular weight from MS




Fig. 4. Migration assay with 3T3 mouse cells